

Importance Of Grazing Management In Improving Water Use Efficiency of Pastures with Tropical Forage Grasses

Juan Andrés Cardoso,

Juan Jiménez, Mauricio Sotelo, Idupulapati M. Rao and Michael Peters

Tropical Forages Program

Alliance of Bioversity International and CIAT

Cali, Colombia


j.a.cardoso@cgiar.org

Joint XXIV International Grassland Congress and XI International Rangeland Congress
23-29 October 2021, Nairobi, Kenya

Importance Of Grazing Management In Improving Water Use Efficiency of Pastures of Tropical Forage Grasses




Why?

- Water use efficiency (WUE) is a plant trait sought to improve drought resistance



Importance Of Grazing Management In Improving Water Use Efficiency of Pastures of Tropical Forage Grasses



How to improve water use efficiency in pastures?



Importance Of Grazing Management In Improving Water Use Efficiency of Pastures of Tropical Forage Grasses

How grazing management can improve water use efficiency (WUE) in pastures?

- Grazing management of pastures planted with tropical forage grasses can improve WUE values if it moderates the process of evapotranspiration




Objective

- To study changes in WUE by simulated grazing under greenhouse conditions in a set of forage grasses commonly used in the tropics (*Cenchrus ciliaris*, *Chloris gayana*, *Megathyrsus maximus*, *Urochloa* spp.) under well-watered and water-limited conditions

Importance Of Grazing Management In Improving Water Use Efficiency of Pastures of Tropical Forage Grasses

Materials and methods

- *Urochloa humidicola* cv. Tully
 - *U. brizantha* cv. Piata
 - *U. hybrid* cv. Mulato II
 - *Megathyrsus maximus* cv. Mombaza
 - *M. maximus* cv. Sabanera
 - *Chloris gayana* ILRI-645,
 - *Cenchrus ciliaris* ILRI-10097
- Two watering conditions (well-watered and water-limited)
 - Three grazing intensities, every 25 days for 100 days



Importance Of Grazing Management In Improving Water Use Efficiency of Pastures of Tropical Forage Grasses

Results

Dry mass forage (g plant ⁻¹)						
Well-watered			Water-limited			
Grazing intensity			Grazing Intensity			
	Low	Moderate	High	Low	Moderate	High
<i>U. humidicola</i> cv. Tully	39.8bc	32.3b	30.3b	22.3b	24.7bc	14.1a
<i>M. maximus</i> cv. Sabanera	83.5g	67.5e	62.2e	30.1bc	43.6d	22.5a
<i>M. maximus</i> cv. Mombaza	99.1h	81.9g	75.5f	29.7bc	47.5e	21.5b
<i>C. gayana</i> ILRI-645	67.7e	53.1d	47.9cd	26.1c	37.0d	25.7c
<i>U. brizantha</i> cv. Piata	55.8d	47.7cd	44.1c	28.4c	36.4d	23.8c
<i>U. hybrid</i> cv. Mulato II	54.5d	50.7d	41.8c	29.9bc	37.1d	25.6bc
<i>C. ciliaris</i> ILRI-10097	33.2b	27.0a	24.5a	19.1ab	24.7bc	18.3ab
Average	60.5e	51.5d	46.6c	26.5c	35.9d	21.6ab

- High grazing intensity resulted in greater forage yield under well-watered conditions, but lower forage yield under water-limited conditions

Leaf rolling scores				Dead leaf dry mass (g plant ⁻¹)		
Water-limited				Water-limited		
Grazing intensity				Grazing Intensity		
	Low	Moderate	High	Low	Moderate	High
<i>U. humidicola</i> cv. Tully	3c	1a	1a	2.6e	0.4ab	0.3a
<i>M. maximus</i> cv. Sabanera	5e	2b	3c	3.2g	0.3a	0.2a
<i>M. maximus</i> cv. Mombaza	5e	2b	3c	5.1f	1.2c	1.0c
<i>C. gayana</i> ILRI-645	5e	2b	2b	2.4e	1.1c	0.5ab
<i>U. brizantha</i> cv. Piata	4d	2b	3c	0.8bc	0.3a	0.3a
<i>U. hybrid</i> cv. Mulato II	4d	2b	3c	1.4cd	0.5ab	0.2a
<i>C. ciliaris</i> ILRI-10097	3c	1a	2b	0.5ab	0.1a	0.1a
Average	4d	2b	2.4b	2.3e	0.6ab	0.4ab

- Plants under low grazing intensity showed greater symptoms of stress under water-limited conditions

Importance Of Grazing Management In Improving Water Use Efficiency of Pastures of Tropical Forage Grasses

Results


	Water Use Efficiency (g plant biomass/liter of water consumed)					
	Well-watered			Water-limited		
	Grazing intensity			Grazing Intensity		
	Low	Moderate	High	Low	Moderate	High
<i>U. humidicola</i> cv. Tully	6.0abc	6.2abc	5.8ab	2.2c	3.2e	2.7d
<i>M. maximus</i> cv. Sabanera	5.3a	5.5a	5.4a	2.0bc	3.0de	1.1a
<i>M. maximus</i> cv. Mombaza	5.5a	5.4a	5.5a	1.5ab	2.5cd	1.4ab
<i>C. gayana</i> ILRI-645	5.9ab	5.9ab	6.0abc	1.5ab	2.5cd	1.0a
<i>U. brizantha</i> cv. Piata	6.3bcd	6.8cd	6.5bcd	2.5cd	3.5ef	1.0a
<i>U. hybrid</i> cv. Mulato II	5.9ab	6.0abc	5.9ab	1.8bc	2.8d	1.5ab
<i>C. ciliaris</i> ILRI-10097	6.7cd	7.1d	7.4d	3.1de	4.1g	3.1de
Average	5.9ab	6.1abc	6.1abc	2.1c	3.1e	1.7b

- Plants under moderate grazing intensity showed greater WUE

Importance Of Grazing Management In Improving Water Use Efficiency of Pastures of Tropical Forage Grasses

Conclusions

- WUE of forages can be improved by managing leaf area size
- Moderate grazing management shows promise for mitigating the impacts of drought events and improve resiliency of pastures planted with tropical grasses.
- Current research is underway to test the aforementioned results under field conditions using remote sensing and Internet of things approaches



Thanks everyone

Juan Andres Cardoso
Juan Jiménez, Mauricio Sotelo, Idupulapati M. Rao and Michael Peters
Tropical Forages Program
Crops for Nutrition and Health
Alliance Bioversity-CIAT
j.a.cardoso@cgiar.org

<https://alliancebioversityciat.org/>
<https://www.tropicalforages.info/text/intro/index.html>
<https://www.tropicalgrasslands.info/index.php/tgft>